

First report of the dog louse fly *Hippobosca longipennis* in Romania

A. D. MIHALCA¹, I. R. PĂSTRĂV¹, A. D. SĂNDOR¹, G. DEAK¹,
C. M. GHERMAN¹, A. SARMAȘI¹ and J. VOTÝPKA^{2,3}

¹Department of Parasitology and Parasitic Diseases, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Cluj-Napoca, Romania, ²Department of Parasitology, Faculty of Science, Charles University, Prague, Czech Republic and ³Biology Centre, Institute of Parasitology, Czech Academy of Sciences, České Budějovice, Czech Republic

Abstract. *Hippobosca longipennis* (Diptera: Hippoboscidae), the dog fly or dog louse fly, is an obligate blood-feeding ectoparasite of wild and domestic carnivores in Africa and the Middle East. Outside its typically known geographic range, *H. longipennis* has been reported occasionally on mainly domestic dogs in Asia and southern Europe, and infrequently in other areas (central Europe and the U.S.A.). This paper presents the first report of *H. longipennis* in Romania and the second record of *Lipoptena fortisetosa* (Diptera: Hippoboscidae), a potentially invasive species. *Hippobosca longipennis* was found on domestic dogs in two regions of the country (northern Romania in Maramures and southwestern Romania in Dobrogea) and on two road-killed wildcats in Maramures. *Lipoptena fortisetosa* was found on domestic dogs in Maramures. In both species identification was based on morphology and confirmed by barcoding of the cytochrome *c* oxidase subunit 1 gene. It is not clear for how long *H. longipennis* has been present in central Europe, nor if it was introduced (via the movement of domestic dogs or import of exotic carnivores) or present historically (Holocene remnants). This paper discusses the possible origins of *H. longipennis* in central Europe as its current distribution in the area is sparse and patchy.

Key words. *Hippobosca longipennis*, *Lipoptena fortisetosa*, dogs, Romania.

Introduction

Hippobosca longipennis, the dog fly or dog louse fly, is an obligate blood-feeding ectoparasite of wild and domestic carnivores including Canidae (jackals, foxes and domestic dogs), Viverridae (mongooses and civets), Hyaenidae (striped and spotted hyenas) and Felidae (cheetahs, servals, leopards, lions and domestic cats) (Bequaert, 1953; Sanborn & Hoogstraal, 1953; Maa, 1969; Lloyd, 2002). It has also been found on incidental hosts such as antelopes and birds, and it can occasionally attack humans (Bequaert, 1953; Maa, 1969). It is widely distributed in Africa and the Middle East (Bequaert, 1953) on both wild and domestic animals. Outside this geographic range, *H. longipennis* has been reported mainly on domestic dogs in warm parts of Asia,

North America and Europe (Bequaert, 1953). In the U.S.A., it was introduced through the import of cheetahs to the San Diego Zoo in California (Keh & Hawthorne, 1977) and later reported in various wild animals in California, Texas, Georgia and Oregon (Lloyd, 2002). In Europe, reports of *H. longipennis* come mainly from the Mediterranean countries and more rarely from central Europe (Bequaert, 1953). Occasional findings of imported cases have also been reported from the U.K. (O'Connor & Sleeman, 1987; Chandler, 1998). *Hippobosca longipennis* is considered to be the main vector of the vector-borne filarial nematode parasite *Acanthocheilonema dracunculoides* (Filarioidea: Onchocercidae) (Nelson, 1963; Anderson, 2000). Although *A. dracunculoides* has not yet been reported in Romania, it has been detected in Europe in red foxes from Italy (Magi *et al.*,

Correspondence: Ioana Raluca Păstrav, Department of Parasitology and Parasitic Diseases, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Calea Mănăștur 3–5, Cluj-Napoca 400372, Romania. Tel.: +40 264 596384; Fax: +40 264 593792; E-mail: ioana.pastrav@usamvcluj.ro

2008), in foxes, lynxes and domestic dogs and cats from Spain (Olmeda-García *et al.*, 1993; Bolio *et al.*, 2004; Millán *et al.*, 2007), and in domestic dogs from Portugal (Alho *et al.*, 2014).

Deer keds of the genus *Lipoptena* are obligate blood-feeding ectoparasites that shed their wings after finding a suitable host. Deer keds are potential vectors of various pathogens, including *Anaplasma*, *Bartonella*, *Rickettsia* and *Trypanosoma*. Whereas *Lipoptena cervi* is a native European species, *Lipoptena fortisetosa* was first identified in Japan in the 1960s (Maa, 1965). This probably Asian species was also noted in Europe for the first time in the 1960s (Chalupský, 1980) and was later recorded in numerous countries.

This study presents the first report of *H. longipennis* and the second record of *L. fortisetosa* in Romania.

Materials and methods

During a routine epidemiologic survey for vector-borne diseases performed in September 2016, 73 domestic dogs (all with outdoor housing and from rural environments) were examined for ectoparasites by visual inspection in several villages in southwestern Maramures County, as follows: Buciumi (47.472055 N, 23.489352 W) ($n = 2$); Curtuiusu Mare (47.427404 N, 23.444610 W) ($n = 4$); Fericea (47.407208 N, 23.387726 W) ($n = 21$); Mesteacăn (47.381775 N, 23.518889 W) ($n = 7$); Valea Chioarului (47.427607 N, 23.483813 W) ($n = 27$), and Vărai (47.387184 N, 23.449287 W) ($n = 12$). Between September 2015 and November 2017, seven road-killed wildcats (*Felis silvestris*) were also collected in the same area as follows: Valea Chioarului ($n = 3$); Buciumi ($n = 1$); Mesteacăn ($n = 1$); Vărai ($n = 1$), and Fericea ($n = 1$). All data for positive animals are shown in Table S1. The sampled dogs were housed outdoors, in improvised shelters, together with other domestic livestock. Additionally, louse fly samples were occasionally found on dogs and collected using fine forceps during another study of dog ticks in the Dobrogea region (see Sándor *et al.*, 2014), in Iazurile (Tulcea County) (45.014040 N, 28.941084 N) during August 2013 and July 2015. These flies were collected from community dogs, which were living ferally and without shelter. The presence of louse flies in this area was occasional and no flies were found on later visits. Another louse fly was collected during a routine examination of a dog for ticks in Pesteră (Constanța County) (44.188671 N, 28.111319 W) in June 2016. Indeed there has been a mistake. The dogs from Maramures county are housed outdoors in improvised shelters, while the dogs from Dobrogea region and Tulcea county were community dogs without shelter. All sampling locations are shown in Fig. 1.

All louse flies were collected alive directly from the respective hosts and placed in 70% ethanol and later examined under a stereomicroscope for morphological identification. Morphological species identification was performed using the available keys (Maa, 1962; Chalupský, 1980) and verified using molecular methods. Two randomly selected ethanol-stored specimens of each morphospecies were washed individually for 1 h in distilled water. Three legs of each specimen tested were removed and DNA extracted using a QIAamp DNA Mini Kit (Qiagen GmbH,

Hilden, Germany) according to the manufacturer's instructions and following the protocol for DNA purification from tissues. The barcoding involved the sequencing of the cytochrome *c* oxidase subunit 1 (*cox1*) gene using universal primers and conditions described by Hebert *et al.* (2003). Amplification reactions were performed in a total volume of 20 μ L with EmeraldAmp GT polymerase chain reaction (PCR) master mix (Takara Bio Europe, Saint-Germain-en-Laye, France). Amplified products were visualized on 1% agarose gel, purified by QIAquick PCR Purification Kit (Qiagen GmbH), and directly sequenced in both directions using amplification primers (ABI Prism BigDye Terminator Cycle Sequencing Ready Reaction Kit; Thermo Fisher Scientific, Inc., Waltham, MA, U.S.A.). The obtained sequences were viewed and aligned using the software GENEIOUS Version 10.0.6. The final sequences were compared with those available in the databases BOLD (<http://www.boldsystems.org/>) and GenBank (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Results

Collections from domestic dogs in Maramures County yielded 14 louse flies, all of which were found in one locality, the village of Fericea, on five dogs. Eleven specimens were morphologically identified as *H. longipennis* (Fig. 2) and three as *L. fortisetosa*. Of wildcats from the same area, two animals were found to be infested with louse flies, both with a single individual identified morphologically as *H. longipennis*. Collections from dogs in the Dobrogea region yielded three louse flies (two in Iazurile, one in Pesteră), all of which were identified as *H. longipennis*. The louse flies on all dogs were found in the fur of the dorsal area of the body (i.e. the back, along the caudal part of the cervical spine and the thoracic area of the spine). The geographic distribution of findings is shown in the map in Fig. 1.

Comparisons of the newly obtained *cox1* gene sequences with those available in the databases (BOLD and GenBank) confirmed (identity 98–99%) the morphological determination of both louse fly species. Two identical sequences of *H. longipennis* represent the same haplotype (GenBank accession no. of the specimen voucher HL_Ro_01: MK405667). However, two sequences of *L. fortisetosa* represent two different haplotypes (MK405668 for specimen voucher LF_Ro_01 and MK405669 for LF_Ro_02).

In the survey in Maramures County, a total of five dogs from four households were found to be infested with *H. longipennis* and/or *L. fortisetosa*. In the first household, two of three dogs were infested. Both were mongrels that lived outdoors and were used for security purposes; they were aged 1 year and 4 years, respectively. The second household had one infected mongrel dog aged 5 years. The last two positive households were situated in close proximity to one another (next-door neighbours) and had one dog each. The dogs were aged 7 years and 10 years, respectively. The samples collected from Dobrogea region originated from two different locations. In Iazurile the same household provided two samples in two different years (in both cases from one young mongrel that was kept outdoors), whereas the sample from Pesteră was collected from one of two mongrels examined, both of which were used as sheep dogs.




Fig. 1. Positive and negative locations of *Hippobosca longipennis* and *Lipoptena fortisetosa* in Romania.



Fig. 2. *Hippobosca longipennis* identified on one of the dogs examined in the present study. [Colour figure can be viewed at wileyonlinelibrary.com].

Discussion

This work represents the first record of the dog louse fly (*H. longipennis*) and the second record of the deer louse fly (*L. fortisetosa*) in Romania. *Lipoptena fortisetosa* has been identified in several European countries, including the Czech Republic (Chalupský, 1980; Barták, 1995), Germany (Schumann & Messner, 1993), Poland (Kowal *et al.*, 2009), Lithuania

(Lithuanian Entomological Society, 2006) and Moldavia (Metelitsa & Veselkin, 1989). This species is certainly more widespread in Europe; however, it is probably overlooked and misidentified as *L. cervi* in many cases. If the Asian origin of *L. fortisetosa* is accepted, its most probable route of introduction to Europe was translocation via its host, probably the sika deer (*Cervus nippon*). However, there is also the possibility that this species has been overlooked for a long time and misidentified

as *L. cervi* or has naturally dispersed throughout Eurasia. In Romania, *L. fortisetosa* was recorded only once, in 2004, in two neighbouring localities (Lung Albina islets) on the lower Danube (Pârnu, 2005). Thus, the present finding represents the second piece of evidence for the presence of the species in Romania and refers to a locality geographically distant from that cited in the previous record. Typical hosts of *L. fortisetosa* are deer, antelopes, cattle, goats and sheep; however, similarly to the present findings, the species has also been captured on dogs (Sokół & Gałęcki, 2017).

Although, *H. longipennis* is a very common parasite of domestic and wild carnivores, particularly in East Africa, North Africa and the Middle East (Bequaert, 1953), the species has been reported occasionally in many other countries, mainly on domestic dogs. Most of the European records originate from regions with a warm Mediterranean climate, such as Spain, Italy including Sicily and Sardinia, Greece, Cyprus and Bulgaria (Bequaert, 1953). Wild carnivores are also important hosts in southern Europe. Lariviere & Calzada (2001) reported an unspecified *Hippobosca* (with high probability of being *H. longipennis*) in genets (*Genetta genetta*) in southern Spain. In another study from southern Spain, Millán *et al.* (2007) found prevalence rates of 31% in the endangered Iberian lynx (*Lynx pardinus*), 26% in red foxes (*Vulpes vulpes*) and 2% in domestic cats. Cordero *et al.* (1994) reported *H. longipennis* in domestic dogs from central Spain (Salamanca) and Portugal.

Records in central Europe are far less common. *Hippobosca longipennis* has been sporadically collected from domestic dogs in Hungary, Ukraine (Crimea), Germany, Poland and Slovakia (Bequaert, 1953; Straka & Majzlan, 2010; Sokół & Gałęcki, 2017). Straka & Majzlan (2010) trapped *H. longipennis* in southern Slovakia, although the first finding dates back to 1953 (Chalupský, 1980). More recently, an extensive study was performed in urban dogs from central Poland and four species of louse fly were identified, with *H. longipennis* being the most dominant (Sokół & Gałęcki, 2017). Furthermore, *H. longipennis* was imported to the U.K. and Ireland with wild carnivores from Africa (O'Connor & Sleeman, 1987; Chandler, 1998). However, there is no evidence of its permanent establishment. Moreover, none of the previous studies of ectoparasites of domestic and wild carnivores in Romania reported the presence of *H. longipennis* (Dumitrache *et al.*, 2011; D'Amico *et al.*, 2017a, 2017b; Foley *et al.*, 2017; Sándor *et al.*, 2017). It should be noted that *Hippobosca equina* has also been reported in dogs from Albania (Danielova, 1960) and Poland (Sokół & Gałęcki, 2017).

Similarly to *L. fortisetosa*, it is not clear how long *H. longipennis* has been present in Europe nor whether it was introduced or present historically, particularly in southern Europe (Bequaert, 1953), as its main African hosts were present in some of these areas until the Holocene (i.e. lions as recently as 1000 BC in Greece) (Kingdon & Hoffmann, 2013). Bequaert (1953) mentioned *H. longipennis* in dogs from ancient Greece (c. 800 BC) and Huchet *et al.* (2013) found a single specimen on the mummy of a dog from ancient Egypt. In central Europe, it is not clear if the reported populations are Holocene remnants or if *H. longipennis* has been introduced through the movement of domestic dogs, resident wild carnivores or importation of African wildlife to zoos, as in the U.K., Ireland

and the U.S.A. (Keh & Hawthorne, 1977; O'Connor & Sleeman, 1987; Chandler, 1998). Other authors have hypothesized that *H. longipennis* is under geographic expansion (Sokół & Gałęcki, 2017) as a result of climate change and increases in mean temperatures. However, there is no direct evidence for such a claim. It is interesting that, as was highlighted by Bequaert (1953), the distributions of most other canine parasites are more or less worldwide, whereas *H. longipennis* has a very patchy distribution outside its typical range which does not seem to be related to climatic factors. The factors determining its patchy distribution in non-Mediterranean regions of Europe remain unknown.

The importance of the presence of *H. longipennis* resides in its vectorial capacity, so far clearly demonstrated only for the filarial nematode *A. dracunculoides* (Nelson, 1963). Although the presence of *A. dracunculoides* in central and eastern Europe has not yet been reported despite systematic surveillance in both domestic and wild carnivores (Ionică *et al.*, 2015, 2016, 2017a, 2017b), the presence of its vectors should raise awareness. However, the louse flies (Hippoboscidae) are not the sole vectors of this parasite as ticks [*Rhipicephalus sanguineus sensu lato* (Ixodida: Ixodidae)] have also been incriminated (Olmeda-García *et al.*, 1993; Olmeda-García & Rodríguez-Rodríguez, 1994; Mihalca *et al.*, 2012; Dantas-Torres & Otranto, 2015). Hence, the importance of this finding, together with the underlying factors determining the distribution of *H. longipennis*, require further investigation.

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Table S1. Records of louse flies collected in Romania, including collection date, locality, coordinates (N, latitude; E, longitude), host species, sex, age and origin.

Acknowledgements

This research was supported by the Executive Agency for Higher Education, Research, Development and Innovation Funding (UEFISCDI), Romania (grant no. PCCDI 57/2018) and the European Regional Development (ERD) Fund (project CePaViP; 16_019/0000759). The authors declare no conflicts of interest.

References

- Alho, A.M., Landum, M., Ferreira, C. *et al.* (2014) Prevalence and seasonal variations of canine dirofilariosis in Portugal. *Veterinary Parasitology*, **206**, 99–105.
- Anderson, R.C. (2000) *Nematode Parasites of Vertebrates: Their Development and Transmission*, 2nd edn. CABI Publishing, Wallingford.
- Barták, M. (1995) Diptera communities of 'Osecká' dump near Duchcov and its environment. *Dipterologica Bohemoslovaca*, **7**, 5–16.
- Bequaert, J.C. (1953) The Hippoboscidae or louse-flies (Diptera) of mammals and birds. Part I. Structure, physiology and natural history. *Entomologica Americana*, **33**, 211–442.

- Bolio, M.E., Montes, A.M., Alonso, F.D. *et al.* (2004) Prevalence of *Dipetalonema dracunculoides* in dogs in Murcia, Spain. *Veterinary Record*, **154**, 726–727.
- Chalupský, J. (1980) Hippoboscidae. *Krevsající mouchy a střechci. Fauna ČSSR, Vol. 22* (ed. by M. Chvála *et al.*), pp. 447–478. Academia, Prague (in Czech).
- Chandler, P. (1998) Checklists of insects of the British Isles (New Series). Part I: Diptera (incorporating a list of Irish Diptera). *Handbook for the Identification of British Insects*, Vol. **12**, p. 234. Royal Entomological Society, London.
- Cordero, M., Castañón, L. & Reguera, A. (1994) *Índice-Catálogo de Zooparásitos Ibéricos, 650*. Secretariado de Publicaciones, Universidad de León, León.
- D'Amico, G., Dumitrache, M.O., Matei, I.A. *et al.* (2017a) Ixodid ticks parasitizing wild carnivores in Romania. *Experimental and Applied Acarology*, **71**, 139–149.
- D'Amico, G., Juránková, J., Tăbăran, F.A. *et al.* (2017b) Occurrence of ticks in the subcutaneous tissue of red foxes, *Vulpes vulpes* in Czech Republic and Romania. *Ticks and Tick-borne Diseases*, **8**, 309–312.
- Danielova, V. (1960) Krovososušie dvukrylye (Fam. Tabanidae, Hippoboscidae, Hypodermatidae) iz sborov èkspedicii ČSAN v Albanii. *Ceskoslovenska Parasitologie*, **7**, 37–40.
- Dantas-Torres, F. & Otranto, D. (2015) Further thoughts on the taxonomy and vector role of *Rhipicephalus sanguineus* group ticks. *Veterinary Parasitology*, **208**, 9–13.
- Dumitrache, M.O., Gherman, C.M., Cozma, V. *et al.* (2011) Hard ticks (Ixodidae) in Romania: surveillance, host associations and possible risks for tick-borne diseases. *Parasitology Research*, **110**, 2067–2070.
- Foley, P., Foley, J., Sándor, A.D. *et al.* (2017) Diversity of flea (Siphonaptera) parasites on red foxes (*Vulpes vulpes*) in Romania. *Journal of Medical Entomology*, **54**, 1243–1250.
- Hebert, P.D.N., Cywinska, A., Ball, S.L. & DeWaard, J.R. (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences*, **270**, 313–321.
- Huchet, J.B., Callou, C., Lichtenberg, R. & Dunand, F. (2013) The dog mummy, the ticks and the louse fly: archaeological report of severe ectoparasitosis in Ancient Egypt. *International Journal of Paleopathology*, **3**, 165–175.
- Ionică, A.M., Matei, I.A., Mircean, V. *et al.* (2015) Current surveys on the prevalence and distribution of *Dirofilaria* spp. and *Acanthocheilonema reconditum* infections in dogs in Romania. *Parasitology Research*, **114**, 975–982.
- Ionică, A.M., Matei, I.A., D'Amico, G. *et al.* (2016) Role of golden jackals (*Canis aureus*) as natural reservoirs of *Dirofilaria* spp. in Romania. *Parasites & Vectors*, **9**, 240.
- Ionică, A.M., Zitra, C., Wimmer, V. *et al.* (2017a) Mosquitoes in the Danube Delta: searching for vectors of filarioid helminths and avian malaria. *Parasites & Vectors*, **10**, 324.
- Ionică, A.M., Matei, I.A., D'Amico, G. *et al.* (2017b) Filarioid infections in wild carnivores: a multispecies survey in Romania. *Parasites & Vectors*, **10**, 332.
- Keh, B. & Hawthorne, R.M. (1977) The introduction and eradication of an exotic ectoparasitic fly, *Hippobosca longipennis* (Diptera: Hippoboscidae), in California. *Journal of Zoo and Wildlife Medicine*, **8**, 19–24.
- Kingdon, J. & Hoffmann, M. (2013) *Mammals of Africa*, Vol. **5**. Bloomsbury Publishing, London.
- Kowal, J., Nosal, P., Rościszewska, M. & Matysek, M. (2009) New records of *Lipoptena fortisetosa* Maa, 1965 (Diptera: Hippoboscidae) in Poland. *Dipteron*, **25**, 27–29.
- Lariviere, S. & Calzada, J. (2001) *Genetta genetta*. *Mammalian Species*, **680**, 1–6.
- Lithuanian Entomological Society (2006) *New and Rare for Lithuania Insect Species: Records and Descriptions*, Vol. **17**. LES, Vilnius.
- Lloyd, J.E. (2002) Louse flies, keds, and related flies (Hippoboscoidea). *Medical and Veterinary Entomology* (ed. by G. Mullen & L. Durden), pp. 349–362. Academic Press, San Diego, CA.
- Maa, T.C. (1962) Notes on the Hippoboscidae (Diptera). *Pacific Insects*, **4**, 583–614.
- Maa, T.C. (1965) A synopsis of the Lipopteninae. *Journal of Medical Entomology*, **2**, 233–248.
- Maa, T.C. (1969) A revised checklist and concise host index on Hippoboscidae (Diptera). *Pacific Insects Monograph*, **20**, 261–299.
- Magi, M., Calderini, P., Gabrielli, S. *et al.* (2008) *Vulpes vulpes*: a possible wild reservoir for zoonotic filariae. *Vector-Borne and Zoonotic Diseases*, **8**, 249–252.
- Magnis, J., Lorentz, S., Guardareo, L. *et al.* (2013) Morphometric analyses of canine blood microfilariae isolated by the Knott's test enables *Dirofilaria immitis* and *D. repens* species-specific and *Acanthocheilonema* (syn. *Dipetalonema*) genus-specific diagnosis. *Parasites & Vectors*, **6**, 48.
- Metelitsa, A.K. & Veselkin, G.A. (1989) Parasitism of the louse fly *Lipoptena fortisetosa* on cattle. *Parazitologiya*, **23**, 276–277.
- Mihalca, A.D., Dumitrache, M.O., Magdas, C. *et al.* (2012) Synopsis of the hard ticks (Acari: Ixodidae) of Romania with update on host associations and geographical distribution. *Experimental and Applied Acarology*, **58**, 183–206.
- Millán, J., Ruiz-Fons, F., Márquez, F.J., Viota, M., López-Bao, J.V. & Paz Martín-Mateo, M. (2007) Ectoparasites of the endangered Iberian lynx *Lynx pardinus* and sympatric wild and domestic carnivores in Spain. *Medical and Veterinary Entomology*, **21**, 248–254.
- Nelson, G.S. (1963) *Dipetalonema dracunculoides* (Gobbold, 1870), from the dog in Kenya: with a note on its development in the louse-fly, *Hippobosca longipennis*. *Journal of Helminthology*, **37**, 236–240.
- O'Connor, J.P. & Sleeman, D. (1987) A review of the Irish Hippoboscidae (Insecta: Diptera). *Irish Naturalists' Journal*, **22**, 236–239.
- Olmeda-García, A.S. & Rodríguez-Rodríguez, J.A. (1994) Stage-specific development of a filarial nematode (*Dipetalonema dracunculoides*) in vector ticks. *Journal of Helminthology*, **68**, 231–235.
- Olmeda-García, A.S., Rodríguez-Rodríguez, J.A. & Rojo-Vázquez, F.A. (1993) Experimental transmission of *Dipetalonema dracunculoides* (Cobbold 1870) by *Rhipicephalus sanguineus* (Latreille 1806). *Veterinary Parasitology*, **47**, 339–342.
- Pârnu, C. (2005) Diptera from the green corridor of the Danube (Romania). *Travaux du Muséum National d'Histoire Naturelle 'Grigore Antipa'*, **48**, 147–176.
- Sanborn, C.C. & Hoogstraal, H. (1953) Some mammals of Yemen and their ectoparasites. *Chicago National History Museum*, **34**, 229–252.
- Sándor, A.D., Dumitrache, M.O., D'Amico, G., Kiss, B.J. & Mihalca, A.D. (2014) *Rhipicephalus rossicus* and not *R. sanguineus* is the dominant tick species of dogs in the wetlands of the Danube Delta, Romania. *Veterinary Parasitology*, **204**, 430–432.

- Sándor, A.D., D'Amico, G., Gherman, C.M., Dumitrache, M.O., Domsa, C. & Mihalca, A.D. (2017) Mesocarnivores and macroparasites: altitude and land-use predict the ticks occurring on red foxes (*Vulpes vulpes*). *Parasites & Vectors*, **10**, 173.
- Schumann, H. & Messner, B. (1993) *Lipoptena fortisetosa* Maa, 1965 new in Germany (Dipt., Hippoboscidae). *Entomologische Nachrichten und Berichte*, **37**, 247–249.
- Sokół, R. & Galecki, R. (2017) Prevalence of keds on city dogs in central Poland. *Medical and Veterinary Entomology*, **31**, 114–116.
- Straka, V. & Majzlan, O. (2010) Dynamics of the flies abundance (Diptera) in the National Nature Reserve Bábsky les near Nitra (South Slovakia). *Rosalia (Nitra)*, **21**, 167–184.

Accepted 11 June 2019

First published online 7 July 2019